

## The effects of dietary glutamine supplementation on growth performance and intestinal morphology of broiler chickens reared under hot conditions

Farhad JAZIDEH<sup>1</sup>, Parviz FARHOOMAND<sup>1</sup>, Mohsen DANESHYAR<sup>1\*</sup>, Gholamreza NAJAFI<sup>2</sup>

<sup>1</sup>Department of Animal Science, Faculty of Agriculture, Urmia University, Urmia, Iran

<sup>2</sup>Department of Veterinary Science, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Received: 17.10.2012 • Accepted: 12.03.2013 • Published Online: 21.04.2014 • Printed: 20.05.2014

**Abstract:** Dietary supplementation effects of 0%, 0.25%, 0.5%, and 1% glutamine were investigated on the performance and development of the gastrointestinal tract in broilers reared under hot conditions. No significant differences were observed between the treatments for feed intake and feed conversion ratio during the starter period, grower period, or entire experiment ( $P > 0.05$ ). During the grower period, only 0.5% glutamine-fed birds had higher body weight gain than others ( $P < 0.05$ ). For the entire period, body weight gain of 0.5% glutamine-fed birds was greater than that of 0.00% and 0.25% glutamine-fed ones ( $P < 0.05$ ). Furthermore, 0.5% glutamine-fed birds showed longer villus height than other birds ( $P < 0.05$ ). In the jejunum, both 0.5% and 1% glutamine consumption caused longer villus height as compared to 0.00% and 0.25% glutamine ( $P < 0.05$ ). Villus height of 0.25% glutamine-fed birds was higher than that of 0.00% glutamine-fed ones ( $P < 0.05$ ). None of the ileum morphological parameters were affected by glutamine supplementation ( $P > 0.05$ ). In conclusion, supplementation of 0.5% glutamine improved the intestinal morphology and body weight gain of broilers under heat stress. Higher dietary glutamine (1%) increased the villus height in the jejunum but did not change the performance.

**Key words:** Broiler chickens, glutamine, heat stress condition, intestinal morphology, performance

### 1. Introduction

Stress can have a profound effect on overall physiology, animal health, and productivity. The gastrointestinal tract is particularly responsive to stressors, which can cause a variety of changes including alteration of the normal protective microbiota (1) and decreased integrity of the intestinal epithelium (2). The nutritional stressors have deleterious effects on the absorptive epithelium of the intestine, resulting in reduction of villus height and crypt depth (3). Less is known about the effect of high environmental temperature on intestinal morphology. High environmental temperature decreases feed consumption, live weight, and feed efficiency in broiler chickens (4). Heat stress (HS) also significantly decreases the broilers' jejunal villus height (5). It causes less crypt depth, mucous area, and villus height of the duodenum and lowers the intestinal length in broiler chickens (6). Furthermore, reduced gut motility (7), changes in intestinal microflora (8), and a depression in the gastrointestinal blood flow have been noticed in broilers under HS conditions (9). Acute HS depresses enterocyte proliferation and affects the expression and activity of brush-border membrane enzymes in young birds (10). Thus, HS exposure may result in impaired digestibility of various essential amino

acids. For example, decreased L-leucine transport has been reported across broilers' intestinal epithelium in *in vitro* HS studies (11). Accelerated replacement of enterocytes requires energy and proteins, which can negatively impact the growth and development of other tissues and organ systems (12).

Glutamine (Gln) is the principal metabolic fuel for small intestine enterocytes, lymphocytes, macrophages, and fibroblasts (13) and is considered as an essential amino acid in some species under inflammatory conditions of infection and injury (14). Many benefits of dietary Gln supplementation have been recognized in humans, rats, and pigs. In rats, Gln stimulated the gut mucosal proliferation (15). In another experiment on rats, dietary Gln (1.5%) maintained the gut integrity, which is important in preventing bacterial infections (16). It has even been shown to prevent intestinal hyperpermeability and bacterial translocation in mice during immunological challenge (17). In weanling pigs, Kitt et al. (18) showed improved feed efficiency by dietary addition of 1% Gln. Some beneficial effects of Gln have been recognized in broiler chickens, whereas no information is available on effects during HS. In an experiment on broiler chickens, Yi et al. (19) noticed ameliorated body weight gain and feed

\* Correspondence: m.daneshyar@urmia.ac.ir

efficiency during the first week after hatching with Gln (1%) under normal temperature conditions.

To date, no research has been conducted on the use of dietary Gln in broiler chickens under HS, and so the present experiment evaluated the effect of Gln on growth performance and intestinal morphology of broilers reared under hot conditions.

## 2. Materials and methods

A total of 200 Ross 308 male broiler chicks of 1 day old were used in this study. The chicks were obtained from a local hatchery, weighed on arrival, and randomly allotted into 4 equal groups with 5 replicates each (10 chicks per replicate). The chicks were housed in a clean, well-ventilated room, previously disinfected with formalin. The room was provided with heaters to adjust the environmental temperature to  $32 \pm 1$  °C and this temperature was maintained for 42 days. Relative humidity was 40%. Feeds and water were supplied ad libitum. Prophylactic measures against the most common infectious diseases were carried out. The chicks were vaccinated against Newcastle and Gumboro diseases as prescribed by a local veterinarian.

The treatment diets were formulated to meet the requirements of broiler chickens according to Ross requirement guidelines (Ross Company). The experimental diets were formulated to be isonitrogenous and isocaloric with 0.0%, 0.25%, 0.5%, and 1.0% Gln, which were identified as ZG (zero Gln), LG (low Gln), MG (medium Gln), and HG (high Gln), respectively. One percent finely ground, air-dried sand was used as an inert filler in the basal diet and different dietary levels of Gln were replaced by sand in the experimental diets (Table 1).

Gln with 98% purity (Serva Co., USA) was used in the current experiment. Average body weight gain (BWG) and feed intake (FI) were determined during the starter period (0 to 21 days of age), grower period (22 to 42 days of age), and whole experimental period (0 to 42 days of age). Moreover, the feed conversion ratio (FCR) was calculated for these periods. On days 21 and 42 of age, 1 bird from each replicate pen (5 per treatment) was killed by decapitation to collect the intestine samples. The intestine was removed and divided into 3 segments of duodenum (from gizzard to entry of the bile and pancreatic ducts), jejunum (from entry of the ducts to Meckel's diverticulum), and ileum (from Meckel's diverticulum to the ileocecal junction). The empty weight of these parts was determined separately. Approximately 5 cm of the middle portion of the duodenum, jejunum, and ileum was then cut, digesta was washed away using normal saline, and samples were fixed in 10% neutral buffered formalin. Following histological fixation, the tissues were processed through a standard alcohol dehydration-xylene sequence and embedded in paraffin. From each segment, 5 sections

of 7  $\mu$ m in thickness were made and they were stained with hematoxylin and eosin and periodic acid-Schiff stain. Morphometric analyses of digital photos of light microscopy were performed by means of Image J analysis software (20). Villus height (VH), villus width (VW), crypt depth (CD), muscular layer thickness (MLT), and goblet cells number (GCN) were determined and the VH to CD ratio (VCR) was calculated.

At the end of the experiment, 1 bird from each replicate was randomly selected and sacrificed. The weight and length of the different intestinal segments (duodenum, jejunum, and ileum) and their relative lengths (intestinal segment lengths/small intestine length  $\times$  100) and weights (organ weight/live body weight  $\times$  100) were calculated afterwards. All the data were submitted to the GLM procedure of the SAS program (21) based on a completely randomized design, with 4 treatments and 5 replicates each. Duncan's multiple range test at the level of  $P < 0.05$  was used to denote statistical significance.

## 3. Results

### 3.1. Performance

Dietary supplementation effects of Gln on BWG, FI, and FCR during the starter, grower, and whole experimental periods are shown in Table 2. No Gln effect was detected for FI and FCR during the starter, grower, or whole experimental periods ( $P > 0.05$ ). Similarly, no significant difference was determined between the treatments for BWG during the starter period ( $P > 0.05$ ). During the grower period and the whole period, BWG was affected by Gln ( $P < 0.05$ ). The BWG of MG-fed birds was greater than that of other birds during the grower period ( $P < 0.05$ ), but there was no significant differences between other treatments. For the whole experiment, the BWG of MG-fed birds was higher than that of ZG- and LG-fed birds ( $P < 0.05$ ). No significant difference was noted between HG birds and other treatments for BWG during the whole period ( $P > 0.05$ ).

### 3.2. Intestinal morphology

The effects of dietary Gln on proportional intestinal part lengths and weights are shown in Table 3. The results of our experiment revealed that Gln supplementation had no effect on proportional intestinal part (duodenum, jejunum, and ileum) lengths (relative to intestine length  $\times$  100) and weights (relative to live body weight  $\times$  100) on days 21 and 42 of age ( $P > 0.05$ ).

The intestinal morphology of broilers on day 21 of age is indicated in Table 4. In the duodenum, MG-fed birds had longer VHs than other birds ( $P < 0.05$ ). Gln consumption had no effects on VW, CD, MLT, GCN, or VH to CD ratio ( $P > 0.05$ ). In the jejunum, both the MG and HG birds had a higher VH than LG and ZG birds ( $P < 0.05$ ). Furthermore, the VH of LG birds was higher than

**Table 1.** Ingredient composition and chemical analysis of the experimental diets (fresh basis).

	Starter diet (0–21 days)				Grower diet (22–42 days)			
	ZG	LG	MG	HG	ZG	LG	MG	HG
Ingredients (%)								
Corn	31.08	31.08	31.08	31.08	31.98	31.98	31.98	31.98
Wheat	20.00	20.00	20.00	20.00	25.00	25.00	25.00	25.00
Soybean oil	3.50	3.50	3.50	3.50	3.95	3.95	3.95	3.95
Soybean meal (44%) protein)	39.68	39.68	39.68	39.68	33.92	33.92	33.92	33.92
Dicalcium phosphate	2.10	2.10	2.10	2.10	2.15	2.15	2.15	2.15
Calcium carbonate	1.10	1.10	1.10	1.10	0.86	0.86	0.86	0.86
DL-methionine	0.38	0.38	0.38	0.38	0.08	0.08	0.08	0.08
L-Lysine	0.29	0.29	0.29	0.29	0.22	0.22	0.22	0.22
Vitamin and mineral premix <sup>1</sup>	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50
Salt	0.37	0.37	0.37	0.37	0.34	0.34	0.34	0.34
Sand	1.00	0.75	0.50	0.00	1.00	0.75	0.50	0.00
L-Glutamine	0.00	0.25	0.50	1.00	0.00	0.25	0.50	1.00
Total	100	100	100	100	100	100	100	100
Calculated analysis								
Metabolizable energy (MJ/kg) (kcal/g)	2.85	2.85	2.85	2.85	2.95	2.95	2.95	2.95
DM (%)	85.16	85.16	85.16	85.16	85.43	85.43	85.43	85.43
CP (%)	22.01	22.01	22.01	22.01	20.00	20.00	20.00	20.00
Calcium (%)	1.00	1.00	1.00	1.00	0.90	0.90	0.90	0.90
Available phosphorus (%)	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45
Sodium (%)	0.16	0.16	0.16	0.16	0.15	0.15	0.15	0.15
Arginine (%)	1.54	1.54	1.54	1.54	1.38	1.38	1.38	1.38
Methionine + cystine (%)	1.07	1.07	1.07	1.07	0.73	0.73	0.73	0.73
Tryptophan (%)	2.13	2.13	2.13	2.13	1.94	1.94	1.94	1.94
Glycine (%)	0.97	0.97	0.97	0.97	0.89	0.89	0.89	0.89
Serine (%)	1.14	1.14	1.14	1.14	1.04	1.04	1.04	1.04
Glycine+ serine (%)	2.11	2.11	2.11	2.11	1.93	1.93	1.93	1.93
Histidine (%)	0.60	0.60	0.60	0.60	0.55	0.55	0.55	0.55
Isoleucine (%)	0.97	0.97	0.97	0.97	0.88	0.88	0.88	0.88
Leucine (%)	1.81	1.81	1.81	1.81	1.66	1.66	1.66	1.66
Lysine (%)	1.43	1.43	1.43	1.43	1.24	1.24	1.24	1.24
Methionine (%)	0.70	0.70	0.70	0.70	0.38	0.38	0.38	0.38
Cystine (%)	0.37	0.37	0.37	0.37	0.35	0.35	0.35	0.35
Phenylalanine (%)	2.13	2.13	2.13	2.13	1.04	1.04	1.04	1.04
Tyrosine (%)	0.98	0.98	0.98	0.98	0.89	0.89	0.89	0.89
Phenylalanine + Tyrosine (%)	2.13	2.13	2.13	2.13	1.94	1.94	1.94	1.94
Threonine (%)	0.85	0.85	0.85	0.85	0.77	0.77	0.77	0.77
Valine (%)	1.08	1.08	1.08	1.08	0.98	0.98	0.98	0.98

<sup>1</sup>Supplied per kilogram of diet: retinol, 9000 IU; cholecalciferol, 2000 IU; tocopherol, 18 IU; cyanocobalamin, 15 mg; riboflavin, 6.6 mg; pantothenate, 10 mg; niacin, 30 mg; choline, 500 mg; biotin, 0.1 mg; thiamine, 1.8 mg; pyridoxine, 3 mg; folic acid, 1 mg; menadione, 2 mg; ethoxyquin, 100 mg; zinc, 50 mg; manganese, 100 mg; copper, 10 mg; iron, 50 mg; iodine, 1 mg; selenium, 0.2 mg. ZG = 0.0% glutamine, LG = 0.25% glutamine, MG = 0.5% glutamine, HG = 1% glutamine.

**Table 2.** The effects of L-glutamine on performance of broiler chickens reared under hot conditions during the starter (0 to 21 day of age), grower (22 to 42 days of age), and whole experimental (0 to 42 days of age) periods.

	Treatments	FI (g)	BWG (g)	FCR
Starter	ZG	1066.08	680.50	1.56
	LG	1038.91	661.40	1.57
	MG	1071.62	694.50	1.54
	HG	1052.39	662.50	1.59
	Pooled SEM	7.38	7.29	0.02
	P-value	0.39	0.36	0.86
Grower	ZG	2879.45	1285.13 <sup>b</sup>	2.21
	LG	2918.04	1339.64 <sup>b</sup>	2.17
	MG	3047.62	1478.05 <sup>a</sup>	2.06
	HG	3004.74	1354.13 <sup>b</sup>	2.22
	Pooled SEM	31.27	23.92	0.03
	P-value	0.21	0.02	0.19
Whole the experimental period	ZG	3945.50	1975.25 <sup>b</sup>	1.98
	LG	3957.00	2001.04 <sup>b</sup>	1.98
	MG	4105.80	2179.41 <sup>a</sup>	1.88
	HG	4061.70	2053.05 <sup>ab</sup>	1.97
	Pooled SEM	36.52	27.89	0.02
	P-value	0.39	0.04	0.33

<sup>a, b</sup>: Mean values with different superscripts within each column and period are significantly different ( $P < 0.05$ ). BWG = Body weight gain, FI = feed intake FCR = feed conversion ratio, ZG = 0.0% glutamine, LG = 0.25% glutamine, MG = 0.5% glutamine, HG = 1% glutamine.

**Table 3.** Effect of dietary L-glutamine supplementation on proportional intestinal part (duodenum, jejunum, and ileum) lengths (relative to intestine length  $\times 100$ ) and weights (relative to live body weight  $\times 100$ ) in broiler chickens on day 21 of age and reared under hot conditions.

	Treatments	ZG	LG	MG	HG	Pooled SEM	P
Duodenum	Weight	1.34	1.52	1.36	1.51	0.05	0.39
	Length	22.01	20.05	21.02	21.22	0.82	0.89
Jejunum	Weight	2.95	2.70	3.18	2.74	0.09	0.17
	Length	44.99	43.88	43.12	43.34	0.56	0.68
Ileum	Weight	2.13	1.46	1.77	1.57	0.12	0.18
	Length	32.99	36.07	35.85	35.44	0.58	0.22

ZG = 0.0% glutamine, LG = 0.25%, glutamine, MG = 0.5% glutamine, HG = 1% glutamine.

**Table 4.** Effect of dietary L-glutamine supplementation on duodenum, jejunum, and ileum morphology in broiler chickens on day 21 of age and reared under hot conditions.

	Treatments	Glutamine supplementation (%)				Pooled SEM	P-value
		ZG	LG	MG	HG		
Duodenum	VH ( $\mu\text{m}$ )	1394.73 <sup>b</sup>	1410.18 <sup>b</sup>	1449.45 <sup>a</sup>	1390.97 <sup>b</sup>	6.84	0.0005
	VW ( $\mu\text{m}$ )	142.44	140.77	140.35	145.78	1.29	0.47
	CD ( $\mu\text{m}$ )	149.96	143.27	149.54	146.62	1.39	0.32
	MLT ( $\mu\text{m}$ )	190.48	185.46	182.12	187.97	1.38	0.17
	GCN ( $\text{mm}^2$ )	7162.86	7125.27	7176.23	7108.56	13.37	0.25
	VH to CD ratio	9.31	9.84	9.71	9.49	0.06	0.21
Jejunum	VH ( $\mu\text{m}$ )	1109.02 <sup>c</sup>	1145.36 <sup>b</sup>	1182.11 <sup>a</sup>	1187.54 <sup>a</sup>	9.51	0.0009
	VW ( $\mu\text{m}$ )	81.87	81.45	81.45	82.71	1.01	0.97
	CD ( $\mu\text{m}$ )	96.49	99.83	105.26	103.17	1.33	0.08
	MLT ( $\mu\text{m}$ )	143.69 <sup>a</sup>	133.25 <sup>b</sup>	142.87 <sup>a</sup>	141.19 <sup>a</sup>	1.34	0.006
	GCN ( $\text{mm}^2$ )	8166.20	8180.40	8276.47	8297.36	23.07	0.08
	VH to CD ratio	11.52	11.49	11.23	11.52	0.12	0.84
Ileum	VH ( $\mu\text{m}$ )	489.55	489.55	508.35	529.65	6.90	0.11
	VW ( $\mu\text{m}$ )	75.61	76.44	76.44	74.77	1.10	0.95
	CD ( $\mu\text{m}$ )	69.34	69.76	68.09	66.83	1.00	0.77
	MLT ( $\mu\text{m}$ )	130.74	127.82	131.16	134.08	1.27	0.42
	GCN ( $\text{mm}^2$ )	10,080.14	10,077.63	10,082.64	10,037.53	19.28	0.85
	VH to CD ratio	7.06	7.05	7.47	7.96	0.16	0.13

<sup>a-c</sup>: Mean values with different superscripts within a row are significantly different ( $P < 0.05$ ).

VH = Villus height, VW = villus wide, CD = crypt depth, MLT= muscular layer thickness, GCN = goblet cell number, VH to CD ratio = villus height to crypt depth ratio, ZG = 0.0% glutamine, LG = 0.25% glutamine, MG = 0.5% glutamine, HG = 1% glutamine.

that of ZG birds ( $P < 0.05$ ). No significant differences were seen between treatments for other variables ( $P > 0.05$ ). In the ileum, VH, VW, CD, MLT, GCN, and VH to CD ratio were not affected by Gln in diet ( $P > 0.05$ ).

#### 4. Discussion

Many investigators presented the negative effects of HS on gut morphology. Mitchell and Carlisle (5) indicated the lower jejunal villus height in broiler chickens under HS. Marchini et al. (6) showed that HS reduces the crypt depth, mucous area, villus height of the duodenum, and even intestinal length in broiler chickens. Hence, the present experiment investigated the possible positive effects of Gln on intestinal development and performance of broiler chickens under HS. As expected, consumption of 0.5% Gln in birds under HS conditions resulted in

longer VHs in the duodenum and jejunum. It has been stated that addition of Gln to broiler diets positively affects the gastrointestinal tract and could increase the villi length in different segments of the small intestine under normal temperature conditions (22–24). In an experiment on turkey poults, Gln supplementation increased the intestinal villus height and decreased crypt depth (25). Fischer da Silva et al. (23) observed an increased number of intestinal villi with 1% Gln in feed-restricted broiler chickens. Soltan (22) indicated similar results with Gln in broiler chickens vaccinated against Newcastle disease. In another consistent experiment, Murakami et al. (24) reported longer villi in the jejunum on days 14 and 41 after hatching in broilers fed diets including 1% Gln and vitamin E. The positive effect of Gln on VH is related to its nutritive importance for villus growth. Gln is an important

amino acid for utilization as an energy source for the development of the mucosa and stimulates intestinal cell proliferation, thus increasing the absorptive surface of the gastrointestinal mucosa and the utilization of nutrients (26).

Villus height is positively related to villus surface area (5). The expansion of surface area that occurs with villus growth has been used to explain the increased absorptive capacity, whereas decreased villus height lowers the absorptive capability of the small intestine (3). Hence, a higher villus height increases the intestinal surface area and consequently nutrient absorption (22), and results in better performance. We found better BWG for 0.5% Gln-fed birds, which may be related to their longer VHs at day 21 of age and therefore improved nutrient utilization. There are some consistent studies regarding the positive effects of dietary Gln on performance of broiler chickens and especially BWG. Although Murakami et al. (24) and Maiorka et al. (27) found no effect of Gln on BWG (days 1–41) of broiler chickens, in agreement with our results, Bartell and Batal (28) observed higher BWG during the third week of age with 1% Gln. Soltan (22) reported improved daily BWG in broilers fed 1% Gln under normal temperature conditions. In the same temperature conditions, Devi Priya et al. (29) showed that 0.5% Gln administration increased the body weight of broilers during the entire experimental period.

Even though 1% Gln caused the same villus lengths as 0.5% Gln supplementation, it did not increase the BWG. The first possible reason for this negative effect on BWG is the role of Gln in ammoniogenesis, which increases the energy requirement for excretion of uric acids. The Gln acts as a precursor for ammoniogenesis in the gut and kidneys (30). In birds, ammonia is excreted in feces in the form of uric acid and is involved in uric acid synthesis (22). The second reason is possibly the production of high ammonium ions in 1% Gln-fed birds. The Gln is converted to  $\alpha$ -ketoglutarate and thus generates ammonium ions ( $\text{NH}_4$ ). Although excretion of ammonium ions helps buffer metabolic acidosis in normal temperatures (31), it negatively affects the blood acid–base balance during HS-induced alkalosis and hence does not improve BWG. It is thought that panting during HS results in respiratory alkalosis and higher blood pH consequently. Thus, high dietary inclusion of Gln (1%) in broiler chickens under HS increases the ammonium excretion and exacerbates the HS-induced alkalosis, and, as consequence, it does not improve the performance as compared to 0.5% Gln.

In conclusion, the results of this study suggest that 0.5% Gln in the diet improves the intestinal morphology and consequently the BWG of broiler chicks under hot conditions. Although a higher level of dietary Gln (1%) increased the villus height in the jejunum, it did not change the performance.

## References

1. Bailey MT, Coe CL. Prenatal stress alters bacterial colonization of the gut in infant monkeys. *J Pediatr Gastr Nutr* 2004; 38: 414–421.
2. Soderholm JD, Yates DA, Gareau MG, Yang PC, Macqueen G, Perdue MH. Neonatal maternal separation predisposes adult rats to colonic barrier dysfunction in response to mild stress. *Am J Physiol Gastrointest Liver Physiol* 2002; 283: G1257–G1263.
3. Yamauchi K, Kamisoyama H, Isshiki Y. Effects of fasting and refeeding on structures of the intestinal villi and epithelial cells in White Leghorn hens. *Br Poult Sci* 1996; 37: 909–921.
4. Debski B, Zalewski W, Gralak MA, Kosla T. Chromium-yeast supplementation of broiler in an industrial farming system. *J Trace Elem Med Bio* 2004; 18: 47–51.
5. Mitchell MA, Carlisle AJ. The effect of chronic exposure to elevated environmental temperature on intestinal morphology and nutrient absorption in the domestic fowl (*Gallus domesticus*). *Comp Biochem Physiol* 1992; 101A: 137–142.
6. Marchini CFP, Silva PL, Nascimento MRBM, Beletti ME, Silva NM, Guimaraes EC. Body weight, intestinal morphometry and cell proliferation of broiler chickens submitted to cyclic heat stress. *Inter J Poult Sci* 2011; 10: 455–460.
7. Tur JA, Rial RV. The effect of temperature and relative humidity on the gastrointestinal motility of young broilers. *Comp Biochem Physiol A* 1985; 80: 481–486.
8. Suzuki K, Harasawa R, Yoshitake Y, Mitsuoka T. Effects of crowding and heat stress on intestinal flora, body weight gain, and feed efficiency of growing rats and chicks. *Nippon Juigaku Zasshi* 1983; 45: 331–338.
9. Wolfenson D. The effect of acclimatization on blood flow and its distribution in normothermic and hyperthermic domestic fowl. *Comp Biochem Physiol A* 1986; 85: 739–742.
10. Gonzalez R, Eequerra S, Leeson. Physiological and metabolic responses of broilers to heat stress-implications for protein and amino acid nutrition. *World Poult Sci J* 2006; 62: 282–295.
11. Dibener JJ, Atwell CA, Ivery FJ. Effect of heat stress on 2-hydroxy-4-(methylthio) butanoic acid and DL-methionine absorption measured in vitro. *Poult Sci* 1992; 71: 1900–1910.
12. Markovic R, Šefer D, Krsticv M, Petrujkic B. Effect of different growth promoters on broiler performance and gut morphology. *Arc Med Vet* 2009; 41: 163–169.
13. Andrews FJ, Griffiths RD. Glutamine: Essential for immune nutrition in the critically ill. *Br J Nutr* 2009; 51: S3–S8.

14. Newsholme P. Why is L-glutamine metabolism important to cells of the immune system in health post-immune, surgery or infection? *J Nutr* 2001; 131: 2515–2522.
15. Inoue Y, Grant JP, Snyder MH. Effect of glutamine supplemented intravenous nutrition on survival after *Escherichia coli*-induced peritonitis. *JPEN J Parenter Enteral Nutr* 1993; 17: 41–46.
16. Naka S. Alanine-glutamine-supplemented total parenteral nutrition improves survival and protein metabolism in rat protracted bacterial peritonitis model. *JPEN J Parenter Enteral Nutr* 1996; 20: 417–423.
17. Adjei AA, Matsumoto Y, Oku T, Hiroi Y, Yatsumoto Y. Dietary arginine and glutamine combination improves survival in septic mice. *Nutr Res* 1994; 14: 1591–1599.
18. Kitt J, Miller PS, Lewis AJ, Fischer RL. Effects of Glutamine on Growth Performance and Small Intestine Villus Height in Weanling Pigs. *Nebraska Swine Reports*. Lincoln, NE, USA: University of Nebraska, London; 2002.
19. Yi GF, Allee GL, Frank JW, Spencer JD, Touchette KJ. Impact of glutamine, menhaden fish meal and spray-dried plasma on the growth and intestinal morphology of broilers. *Poult Sci* 2001; 80 (Suppl. 1): 201.
20. Abràmoff MD, Magelhaes PJ, Ram SJ. Image processing with Image J. *Biophoton Int* 2004; 11: 36–42.
21. SAS Institute. SAS Users Guide: Statistics. Cary, NC, USA: SAS Institute Inc.; 2002.
22. Soltan MA. Influence of dietary glutamine supplementation on growth performance, small intestinal morphology, immune response and some blood parameters of broiler chickens. *Int J Poult Sci* 2009; 8: 60–68.
23. Fischer da Silva AV, Majorca A, Borges SA, Santin E, Boleli IC, Macari M. Surface area of the tip of the enterocytes in small intestine mucosa of broilers submitted to early feed restriction and supplemented with glutamine. *Int J Poult Sci* 2007; 6: 31–35.
24. Murakami AE, Sakamoto MI, Natali MRM, Souza LMG, Farnco JRG. Supplementation of glutamine and vitamin E on the morphometry of the intestinal mucosa in broiler chickens. *Poult Sci* 2007; 86: 488–495.
25. Yi GF, Allee GL, Spencer JD, Frank JW, Gaines AM. Impact of glutamine, menhaden fish meal and spray-dried plasma on the growth and intestinal morphology of turkey poults. *Poult Sci* 2001; 80 (Suppl. 1): 201.
26. Samli HE, Senkoylu N, Koc F, Kanter M, Agma A. Effects of *Enterococcus faecium* and dried whey on broiler performance, gut histomorphology and microbiota. *Arch Anim Nutr* 2007; 61: 42–49.
27. Maiorka A, Silva AVF, Santin E, Borges SA, Boleli IC, Macari M. Influencia da suplementação de glutamina sobre o desempenho e o desenvolvimento de vilos e criptas do intestino delgado de frangos. *Arq Bras Med Vet Zootec* 2000; 52: 487–490 (in Portuguese).
28. Bartell SM, Batal AB. The effect of supplemental glutamine on growth performance, development of the gastrointestinal tract, and humoral immune response of broilers. *Poult Sci* 2007; 86: 1940–1947.
29. Devi Priya K, Selvaraj P, Nanjappan K, Jayachandran S, Visha P. Oral supplementation of putrescine and L-glutamine on the growth performance, immunity, intestinal enzymes in the broilers chickens. *Tamil Nadu J Vet Anim Sci* 2010; 6: 250–254.
30. Tapiero H, Mathé G, Couvreur P, Tew KD. Free amino acids in human health and pathologies - II. Glutamine and glutamate. *Biomed Pharmacother* 2002; 56: 446–457.
31. Chasiotis D, Hultman E, Sahin K. Acidotic depression of cyclic AMP accumulation and phosphorylase b to a transformation in skeletal muscle of man. *J Physiol* 1983; 335: 197–204.