
Cell Membranes and Free Radical Research

Volume4, Number1, October 2013

[CONTENTS]

- 221 **Oxidative Stress and Vascular Function**
Andreas Daiber, Michael Mader, Paul Stamm, Elena Zinßius, Swenja Kröller-Schön, Matthias Oelze, Thomas Münzel
- 232 **Effects of beer, Hops (*Humulus lupulus*) on total antioxidant capacity in plasma of stressed subjects**
Lourdes Franco, Rafael Bravo, Carmen Galan, Cristina Sanchez, Ana Beatriz Rodriguez, Carmen Barrigs, Javier Cubero
- 236 **Musa Sapiantum attenuates total antioxidant and lipid profile values in rats with indomethacin-induced gastric ulceration** By Akinlolu, Abdulazeez Adelaja, Ghazali, Olaide Kamal, Salau, Bamidele Adewale, Otulana, Jubril Olatoye, Omowunmi, Mobolaji

Cell Membranes and Free Radical Research

Volum 4, Number 1, October 2013

ISSN Numbers: 1308-4178 (On-line), 1308-416X

Indexing: Google Scholar, Index Copernicus, Chemical Abstracts, Scopus (Elsevier),
EBSCOhost Research Database

EDITOR

Editor in Chief

Mustafa Nazıroğlu, Department of Biophysics,
Medical Faculty, Suleyman Demirel University, Isparta,
Turkey.

Phone: +90 246 211 37 08. Fax: +90 246 237 11 65

E-mail: mustafanaziroglu@sdu.edu.tr

Managing Editor

A. Cihangir Uğuz, Department of Biophysics, Medical
Faculty, Suleyman Demirel University, Isparta, Turkey.

E-mail: biophysics@sdu.edu.tr

EDITORIAL BOARD

Cell Membranes, Ion Channels and Calcium Signaling

Alexei Tepikin, The Physiological Laboratory, University of
Liverpool, Liverpool, UK

Andreas Lückhoff, Institute of Physiology, Medical Faculty,
RWTH-Aachen University, Germany

Andreas Daiber, 2nd Medical Clinic, Molecular Cardiology,
Medical Center of the Johannes Gutenberg University,
Mainz, Germany

Giorgio Aicardi, Department of Human and General
Physiology, University of Bologna, Italy.

Gemma A. Figtree, North Shore Heart Research Group
Kolling Institute of Medical Research University of Sydney
and Royal North Shore Hospital
Sydney, AUSTRALIA.

Jose Antonio Pariente, Department of Physiology,
University of Extremadura, Badajoz, Spain.

James W. Putney, Jr. Laboratory of Signal Transduction,
NIEHS, NC, USA.

Martyn Mahaut Smith, Department of Cell Physiology and
Pharmacology, University of Leicester, Leicester, UK.

Stephan M. Huber, Department of Radiation Oncology,
Eberhard - Karls University Tübingen, Germany

Enzymatic Antioxidants

Michael Davies, Deputy Director, The Heart Research
Institute, Sydney, Australia.

Süleyman Kaplan, Department of Histology and
Embryology, Medical Faculty, Samsun, Turkey

Xingen G. Lei, Molecular Nutrition, Department of Animal
Science, Cornell University, Ithaca, NY, USA

Ozcan Erel, Department of Biochemistry, Medical Faculty,
Yıldırım Beyazıt University.

Nonenzymatic Antioxidants, Nutrition and Melatonin

Ana B. Rodriguez Moratinos, Department of Physiology,
University of Extremadura, Badajoz, Spain.

Cem Ekmekcioglu, Department of Physiology, Faculty of
Medical University of Vienna, Austria.

Peter J. Butterworth, Nutritional Sciences Division, King's
College London, London, UK

AIM AND SCOPES

Cell Membranes and Free Radical Research is a print and
online journal that publishes original research articles,
reviews and short reviews on the molecular basis of
biophysical, physiological and pharmacological processes
that regulate cellular function, and the control or
alteration of these processes by the action of receptors,
neurotransmitters, second messengers, cation, anions,
drugs or disease.

Areas of particular interest are four topics. They are;

A- Ion Channels (Na⁺ - K⁺ Channels, Cl⁻ channels, Ca²⁺
channels, ADP-Ribose and metabolism of NAD⁺, Patch-
Clamp applications)

B- Oxidative Stress (Antioxidant vitamins, antioxidant
enzymes, metabolism of nitric oxide, oxidative stress,
biophysics, biochemistry and physiology of free oxygen
radicals)

C- Interaction Between Oxidative Stress and Ion Channels

(Effects of the oxidative stress on the activation of the
voltage sensitive cation channels, effect of ADP-Ribose
and NAD⁺ on activation of the cation channels which
are sensitive to voltage, effect of the oxidative stress on
activation of the TRP channels)

D- Gene and Oxidative Stress (Gene abnormalities.
Interaction between gene and free radicals. Gene
anomalies and iron. Role of radiation and cancer on
gene polymorphism)

READERSHIP

Biophysics
Biochemistry
Biology
Biomedical Engineering
Pharmacology
Physiology
Genetics
Cardiology
Neurology
Oncology
Psychiatry
Neuroscience

Keywords

Ion channels, cell biochemistry, biophysics, calcium
signaling, cellular function, cellular physiology,
metabolism, apoptosis, lipid peroxidation, nitric oxide
synthase, ageing, antioxidants, neuropathy.

Effects of beer, Hops (*Humulus lupulus*) on total antioxidant capacity in plasma of stressed subjects

Lourdes Franco¹, Rafael Bravo¹, Carmen Galan¹, Cristina Sanchez¹, Ana Beatriz Rodriguez¹, Carmen Barrigs¹, Javier Cubero^{1,2}.

¹ Neuroimmunophysiology and Chrononutrition Research Group, Department of Physiology. University of Extremadura, Badajoz, Spain

² Laboratory of Health Education, Experimental Sciences Education Area. University of Extremadura, Badajoz, Spain.

List of abbreviations

?????

Corresponding Address

Javier Cubero Juárez
Laboratory of Health Education
Area of Experimental Sciences Education
University of Extremadura
Av. Elvas s/n 06006 Badajoz, SPAIN
Phone & Fax: +34 924 286576
E-mail: jcubero@unex.es

Acknowledgments

This research has been supported by Gobierno de Extremadura, (Fondos FEDER), PII-Acción VII-UEx and Centro de Información Cerveza y Salud (CICS).

Abstract

The fermented drinks of low graduation like beer has many healthy properties such as low-calories, protein compounds, minerals and trace elements, vitamins A, D and E, carbon dioxide and polyphenols. Beer is the only drink which contains Hops (*Humulus lupulus*) and this plant produces a greater and better antioxidant effect. The objective of our experiment was measured the *Antioxidant Capacity* after consumed a non-alcoholic beer. The trial was conducted in a stressed population of 17 female healthy volunteers who drank beer in dinner during two weeks. We analyzed the *Antioxidant Capacity* in urine. The results showed that the consumption of non-alcoholic beer at dinner increases the *Antioxidant Capacity*. In conclusion, our assayed population which ingested a non-alcoholic beer increased the antioxidant levels.

Keywords

Antioxidant, hops, beer, resveratrol and xanthohumol.

Introduction

The source for the majority of life is photosynthesis which turns sun energy into redox energy (Demmig-Adams, 2002). Plants have high concentration of active antioxidants which work like a secondary metabolite, for example, alcohols and polyphenols like xanthohumol and resveratrol, and also carotenoids, glutation, ascorbic acid, antioxidant enzymes, which help to prevent dangerous damage as oxidative stress to cell components (Demmig-Adams, 2002; Benzie, 2003; and Pinto, 2012).

Beer is a fermented drink has many nutritional properties, and contains polyphenols and low levels of graduation (0.1-5%) of alcohol, which generate in moderate consumption an effect healthy (Sánchez, 2010; Arranz, 2012). Beer is the only drink that contains Hops (*Humulus lupulus*) which give the typical bitter flavor. The polyphenols from the hops and the beer: xanthohumol and resveratrol generate in this beverage the antioxidant properties and may reach 0,148 mmol/100 g depending on the type of beer (Bente, 2011).

Flavonoids are a group of natural substances found in the plant kingdom. Have discovered more than 4000 different species of flavonoids, found in fruits, vegetables, seeds, stems, flowers, thus being constituents important in the human diet. His basic structure is: 2 rings benzenes bonded through a ring pyrone (Bravo, 1998). The chemical structure of the phenolic compounds, is conferred by their ability to act as free radical scavengers (Heim, 2002). Beer contains a significant amount of flavonoids. Researchers from King's College London have published an investigation into the presence of flavonoids in various types of beverages and potential role as an antioxidant ferulic acid is found in low alcohol beers (Bourne, 2000). In another study of Oregon State University has shown that hops flavonoids dependent monooxygenases can inhibit cytochrome P-450, suggesting a potential role in modulating the activation of carcinogens (Henderson, 2000). The same group of researchers has confirmed that chalcones and flavanones are inducers of quinone reductase and may have a preventive role in the progression of hepatomas (Miranda, 2000).

Xanthohumol is one of the major flavonoids present in beer from Hops. This biomolecule is used to preserve beer as well as to give the characteristic aroma and flavor to beer (Pinto, 2012). The mechanism of action of xanthohumol has not yet been fully elucidated. Although the evidence for the beneficial effect of xanthohumol is promising, most studies were carried out in vitro. Recently, it has been reported that supplementing the drinking water of rats with xanthohumol prevents

against heterocyclic aromatic mutagens which induce preneoplastic lesions and DNA damage in the liver and colon (Ferk, 2010).

Resveratrol (3, 4, 5-trihydroxy-trans-stilbene) has been found in some hop varieties (Chiva-Blanch, 2010). It is a natural nonflavonoid polyphenolic found in the skin of red grapes (Shakibaei, 2009). Its antioxidant capacity is based in changes in low-density lipoproteins (LDL) properties by oxidation of polyunsaturated fatty acids (PUFA) is believed to play a major role in atherosclerosis. The oxidation affects the protein moiety (Apo B) of LDL particles impairing their catabolism by the regulated apo B/E receptor system. Therefore, the protective role of foods rich in phenolic compounds has been attributed to their antioxidant properties (Rice-Evans, 1997) were the first to demonstrate that trans-resveratrol added to human LDL, reduced the copper-catalyzed oxidation (Frankel 1993). Other benefits of resveratrol are the prevention of the inflammation and cellular senescence (Chung, 2012).

For all theses scientific arguments the objective of our experiment was measured the *Antioxidant Capacity* after consumed a non-alcoholic beer, in a stressed population.

Materials and methods

Subjects

The trial was conducted in a population of 17 female volunteers (auxiliary, nurses and doctors) belonging to the Extremadura Health Service (SES) Badajoz (Spain), which work in rotating shifts and / or night shift, and with an elevated level of work stress and / or mood (assessed by a questionnaire of stress). These women were healthy (40.9±10.5 years), with normal weight and not under any prescription that might influence the study objectives. This project was approved by the Ethics Committee of the University of Extremadura.

Antioxidant Analysis

We chose a longitudinal intervention design in which each subject was his own control. The experimental period was 3 weeks, being the first 7 days the (Week Control), in which the individuals did not ingests any beer at dinner, until the second week (Week 1), in which the individuals ingested 330 mL of non-alcoholic beer during dinner, and the third week (Week 2) is the second week that consumed beer. Urine samples were collected at the end of each of the three weeks of the study. The *Antioxidant Capacity* was analyzed by the improved spectrophotometric "TEAC" method (*Trolox Equivalent Antioxidant Capacity*) (Cubero, 2009). This analysis allows us to calculate the

percent inhibition of ABTS-+ radical cation by Trolox, a water soluble analogue of alpha-tocopherol (vitamin E), which is used as a standard antioxidant. The analysis was carried out on a TECAN plate reader M200 (Männedorf / Switzerland). The wavelength for the study was 730 nm.

Statistical Analysis

After confirming that the population belonged to the Normal, the results were analyzed by *t-Student* test for comparisons between the two groups. We used software GraphPad Prism 5.

Results

Results obtained in a population with high levels of stress (Table 1), showed that the consumption of non-alcoholic beer at dinner increases the antioxidant levels (0.02 mg Trolox) measured in urine.

We observed a growing of *Antioxidant Capacity* in Treatment Group after drinking a non-alcoholic beer for two weeks (0.15±0.03 mg Trolox), versus the Control group without drinking a beer at dinner (0.13±0.01 mg Trolox, Figure 1).

Discussion

It is demonstrated that of free radicals with drinking non-alcoholic beer at dinner is very important for people who work at night because in those hours of sleep, our system acts against free radicals produced in our diurnal life, in addition to fighting viruses and bacteria. Besides, the components of beer are very important of treated oxidative damage, for example, Hops have been used to treat a variety of pathological processes including rheumatism, inflammation and climacteric complaints (Zanoli, 2008). Thus the polyphenol xanthohumol prevents the acute hepatic injury induced by carbon tetrachloride (CCl₄) in rats (Pinto, 2012); respecting to polyphenols many of the biological effects of them have been attributed to their antioxidant properties, either through their reducing capacities per se or through their possible influences on intracellular redox status (Angeloni, 2012).

Table 1. Demographic values of the stressed population (n=17).

Parameters	Mean X±SD
Age (years)	40.90±10.50
Weight (kg)	62.40±11.50
Height (m)	1.61±0.07
CMI (kg/m ²)	24.40±5.30

We report that treatment with non-alcoholic beer shows antioxidant effect of Hops. Our data suggest that the protective properties of non-alcoholic beer are based on the antioxidant effect of the components of Hops: resveratrol and xanthohumol. Our *in vivo* results agree with previous *in vivo* studies (Pinto 2012) and *in vitro* studies reporting the efficiency of polyphenols (Rodriguez 2001).

There were found similar results with the non-alcoholic beer and this *antioxidant ability* (Gonzalez, 2001), by Garrido, (2010) and Bravo, (2012) using cherries and cereals enriched in tryptophan respectively which were supplied to people in dinner.

Referent our trail in stressed population, this *Antioxidant Capacity* will be more higher with higher doses (660 ml) and more time (30 days) of ingest of non alcoholic beer showing the indications of the investigations by Codoñer-Franch, 2012 and Arnao, 2001. For this reason we recommend to increase the ingestion the nonalcoholic beer with this higher dose and more duration.

Apart from that polyphenols show antimicrobial activity, being primarily responsible for the presence of gallic acid. This acid is a natural defense mechanism in many fruits against the development of microbial and fungal diseases (Chung, 1993).

Finally, highlight that the bitter acids components of the Hops modulate on the amino acid GABA (main inhibitory neurotransmitter in the central nervous system), at the same time as myrcenol a terpenoid and xanthohumol in beer from Hops improve its sedative properties (Franco, 2012a and Franco, 2012b).




Figure 1. Comparison of *Antioxidant Capacity* (mg of Trolox) between the Control Group and Group of two weeks of Treatment with non-alcoholic beer.

Conclusion

Drinking a non-alcoholic beer could increase *Antioxidant Capacity* which was measured in the urine from stressed population.

Acknowledgments

This research has been supported by Government of Extremadura Spain, (FEDER), PII-Acción VII-UEx and *Centro de Información Cerveza y Salud (CICS)* Spain.

References

- Angeloni C, Pirola L, Vauzour D, Maraldi T. 2012. Dietary polyphenols and their effects on cell biochemistry and pathophysiology. *Oxidative Medicine and Cellular Longevity*. Hindawi Publishing Corporation.
- Arnao MB, Cano A, Alcolea JF, Acosta M. 2001. Stimulation of free radical-quenching activity of leaf pigment extract. *Phytochem Anal* 12:138-143.
- Arranz S, Chiva-Blanch G, Valderas-Martínez P, Medina-Remón A, Lamuela-Raventós RM and Estruch R. 2012. Wine, Beer, Alcohol and Polyphenols on Cardiovascular Disease and Cancer. *Nutrients* 4:759-781.
- Halvorsen B, Carlsen M, Phillips K, Bøhn S, Holte K, Jacobs D, Blomhoff R. 2006. Content of redox-active compounds (ie, antioxidants) in foods consumed in the United States. *Am J Clin Nutri*. 84:95-135.
- Benzie IF. Evolution of dietary antioxidants. *Comp Biochem Physiol* 2003, 136:113-26.
- Bourne L, Paganga G, Baxter D, Hughes P, Rice-Evans C. 2000. Absorption of ferulic acid from low alcohol beer. *Free Radical Research* 32: 273-280.
- Bravo L. 1996. Polyphenols: Chemistry, dietary sources, metabolism and nutritional significance. *Nutr Rev* 56: 317-333.
- Bravo R, Matito S, Cubero J, Paredes SD, Franco L, Rivero M, Rodríguez AB, Barriga C. 2012. Tryptophan-enriched cereal intake improves nocturnal sleep, melatonin, serotonin, and total antioxidant capacity levels and mood in elderly humans. *AGE* 7(7):e37290
- Chiva-Blanch G, Urpi-Sarda M, Rotchés-Ribalta M, Zamora-Ros R, Llorach R, Lamuela-Raventós R, Estruch R, Andrés-Lacueva C. 2011. Determination of resveratrol and piceid in beer matrices by solid-phase extraction and liquid chromatography-tandem mass spectrometry. *J of Chromat A* 1218: 698-705.
- Chung KT; Stevens SE. *Environ. Toxicol. Chem.* 1993, 12: 2121-2132.
- Chung S, Yao H, Caito S, Hwang JW. 2012. Arunachalam G, Rahman I. Regulation of SIRT1 in cellular functions: role of polyphenols *Arch. Biochem. Biophys.* 501: 79-90.
- Codoñer-Franch P, Hernández-Aguilar MT, Navarro-Ruiz A, López-Jaén AB, Borja-Herrero C, Valls-Bellés V. 2012. Diet Supplementation During Early Lactation with Non-alcoholic Beer Increases the Antioxidant Properties of Breastmilk and Decreases the Oxidative Damage in Breastfeeding Mothers. *Breastfeed Med*.
- Cubero J, Sánchez CL, Bravo R, Sánchez J, Rodríguez AB, Rivero M, Barriga C. 2009. Analysis of the Antioxidant Activity in Human Milk, day vs. Night. *Cell Membranes and free radical research*. 3:100-101.
- Demmig-Adams B, Adams W. 2002. Antioxidants in photosynthesis and human nutrition. *Science*. 298:2149-2153.
- Ferk F, Huber WW, Filipic M, Bichler J, Haslinger E, Misik M, Nersesyanyan A, Grasl-Kraupp B, Zegura B, Knasmüller S. 2010. Xanthohumol, a prenylated flavonoid contained in beer, prevents the induction of preneoplastic lesions and DNA damage in liver and colon induced by the heterocyclic aromatic amine amino-3-methyl-imidazo[4,5-f] quinoline (IQ). *Mutat. Res.* 691: 17-22.
- Franco L, Sánchez C, Bravo R, Rodríguez AB, Barriga C, Romero E, Cubero J. 2012 b. The sedative effects of non-alcoholic beer in healthy female nurses. *PLoS* 7: e37290.
- Franco L, Sánchez CL, Bravo R, Barriga C, Cubero J. 2012. The sedative effects of hops (a component of beer), on activity/rest rhythms. *Acta Physiol Hung.* 20:32-28.
- Frankel EN, Waterhouse AL, Kinsella JE. 1993. Inhibition of human LDL oxidation by resveratrol *Lancet*. 341: 1103-1104.
- Garrido M, Paredes SD, Cubero J, Lozano M, Toribio-Delgado AF, Muñoz JL, Reiter R, Barriga C, Rodríguez AB. 2010. Jerte Valley Cherry-Enriched Diets Improve Nocturnal Rest and Increase 6-Sulfatoxymelatonin and Total Antioxidant Capacity in the Urine of Middle-Aged and Elderly Humans. *J Gerontol A Biol Sci Med Sci.* 65:909-914.
- Heim KE, Tagliaferro AR, Bobilya DJ. 2002. Flavonoid antioxidants: chemistry, metabolism and structure-activity relationships. *J Nutr Biochem* 13: 572-584.
- Henderson MC, Miranda CL, Stevens JF, Deinzer ML, Buhler DR. 2000. In vitro inhibition of human P450 enzymes by prenylated flavonoids from hops. *Humulus lupulus*. *Xenobiotica* 30: 235-251.
- Miranda CL, Aponso GL, Stevens JF, Deinzer ML, Buhler DR. 2000. Prenylated chalcones and flavanones as inducers of quinone reductases in mouse Hepa 1c1c7 cells. *Cancer Letts* 149: 21-29.
- Pinto C, Duque AL, Rodríguez-Galdón B, Cestero JJ, Macías P. 2012. Xanthohumol prevents carbon tetrachloride-induced acute liver injury in rats. *Food Chem Toxicol.* 50:3405-3412.
- Rice-Evans CA, Miller J, Paganga G. 1997. *Trends Plant Sci* 2: 152-159.
- Rodríguez RJ, Miranda CL, Stevens JF, Deinzer ML, Buhler DR. 2001. Influence of prenylated and non-prenylated flavonoids on the liver microsomal lipid peroxidation and oxidative injury in rats hepatocytes. *Food Chem Toxicol* 39:437-445.
- Shakibaei KB, Harikumar BB. 2009. Aggarwal Resveratrol addiction: to die or not to die. *Mol Nutr Food Res* 53: 115-128
- Zanolli P, Zavatti M. 2008. Pharmacognostic and pharmacological profile of *Humulus lupulus* L. *J. Ethnopharmacol* 116: 383-396.

Cell Membranes and Free Radical Research

SUBSCRIPTION ORDER FORM

I would like to subscribe to **Cell Membranes and Free Radical Research**
(ISSN Numbers : 1308-4178 (online) 1308-416X). All prices in Turkish Liras
2012, Vols. (4 issues)

Intitutional Rates:

Print Only: 30 TL per issue + postage 100 TL per year

Credit Card _____ Expiration Date _____

Print Name _____ Signature _____

Receiving Address

Name _____

Department/Title _____

Street _____

City _____ State/Province _____ Post/Zip Code _____ Country _____

Tel _____ Fax _____ E-mail _____

Billing Address Check here if same as above

Name _____

Department/Title _____

Street _____

Cell Membranes and Free Radical Research

LIBRARY ROUTING CARD

To (Librarian) : _____ Date _____

From: _____ Department _____

I have reviewed **Cell Membranes and Free Radical Research**

(ISSN Numbers : 1308-4178 (online) 1308-416X) and suggest that you subscription with:
2012, Vols. (4 issues) All prices in Turkish Liras.

Intitutional Rates:

Print Only: 30 TL + postage 100 TL per year

Süleyman Demirel Üniversitesi Tıp Fakültesi, Isparta / TURKEY

Tel:+ 90 246 211 37 08 Fax:+ 90 246 237 11 65 EMAIL: mustafanaziroglu@sdu.edu.tr

COPYRIGHT FORM

Cell Membranes and Free Radical Research

Date _____ Contributor Name _____

Contributer Address _____

Manuscript Number (if Know) _____

Re: Manuscript entitled _____

“For publication in Cell Membranes and Free Radical Research published by Society of Cell Membranes and Free Oxygen Radicals “

Submission of a manuscript implies:

- that the work described has not been published before (except in the form of an abstract or as part of a published lecture, review or thesis);
- that it is not under consideration for publication elsewhere;
- that its publication has been approved by all co-authors, if any, as well as by the responsible authorities at the institute where the work has been carried out;
- that, if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to this publisher;
- that the manuscript will not be published elsewhere in any language without the consent of the copyright holders;
- that written permission of the copyright holder is obtained by the authors for material used from other copyrighted sources, and that any costs associated with obtaining this permission are the authors' responsibility. Copyright notice: The contributor and the company/employer agree that all copies the final published version of the contribution or any part thereof distributed or posted by them in print or electronic format as permitted herein will include the notice of copyright as stipulated in the journal and a full citation to the journal as published by Cell Membranes and Free radical Society, Isparta, Turkey.

CHECK ONE BOX

Contributor owned work: Contributor's signature _____ Date _____
Type or print name and title _____
Co-Contributor's signature _____ Date _____
Type or print name and title _____

Company-Institution owned work _____
Company-Institution (Employer for here) _____ Date _____
Authorized signature _____ Date _____

U.S. Government work _____

U.K. Government work _____

Other Government work _____